arXiv:1202.2830v2 [gr-qc] 22 Feb 2012

UFES 2012

Thermodynamics of phantom Reissner-Nordstrom-AdS black hole

Deborah F. Jardim $^{(c)}$[1] , Manuel E. Rodrigues $^{(a)}$[2] and Stéphane J. M. Houndjo $^{(b)}$[3]

(a) Universidade Federal do Espírito Santo
Centro de Ciências Exatas - Departamento de Física
Av. Fernando Ferrari s/n - Campus de Goiabeiras
CEP29075-910 - Vitória/ES, Brazil
(b) Institut de Mathématiques et de Sciences Physiques (IMSP)
01 BP 613 Porto-Novo, Bénin
(c) Universidade Federal dos Vales do Jequitinhonha e Mucuri, ICTM
Rua do Cruzeiro, 01, Jardim São Paulo
CEP39803-371 - Teofilo Otoni, MG - Brazil

Abstract

We obtain a new solution of the Einstein-anti-Maxwell theory with cosmological constant, called anti-Reissner-Nordstrom-(A)de Sitter (anti-RN-(A)dS) solution. The basic properties of this solution were reviewed. Its thermodynamics is consistently established, with the extreme cases and phase transitions, making the analysis through two methods, the usual and that of Geometrothermodynamics. The analysis by Geometrothermodynamics does not provide us a result in agreement with the usual method, and by the specific heat. We establish local and global thermodynamic stability of anti-RN-AdS solution through the specific heat and the canonical and grand-canonical ensembles.

Pacs numbers: 04.70.-s; 04.20.Jb; 04.70.Dy.

1 Introduction

The discovery made by Hawking [1], that black holes radiate particles by gravitational interaction near the event horizon, being interpreted like a temperature of the black hole, allowed numerous conjectures for various thermodynamic systems for a variety of black holes solutions in General Relativity,

---

[1] E-mail address: dfjardim@gmail.com
[2] E-mail address: esialg@gmail.com
[3] E-mail address: sthoundjo@yahoo.fr

and other modifications of gravity. This analysis of thermodynamic systems of black holes, as well as the usual thermodynamics, has its base in four postulates, called zero law and three other laws, by analogy to the usual system of thermodynamics [2, 3]. This becomes quite interesting, since each new black hole solution can possess a different thermodynamic system, but which is ruled by laws, or fundamental principles, which are often up to universal, as in usual thermodynamics. The resulting properties of these principles can give us good indications on the local or global stability of these systems, so a good criterion for finding a solution that can be physically acceptable or not.

The thermodynamic properties of a black hole solution can be analysed in many ways. One of them is the usual study made by Davies [4]. But there are other methods well established both physically and mathematically, which are similarly used to study the thermodynamic properties, such as that of the Geometrothermodynamics [5] and that of Hamiltonian thermodynamics [6]. Our main goal here is to use the first two methods in parallel to facilitate comparison of results and their analysis. Because these methods can give us a good notion of the thermodynamic stability of the system, we will study the properties of a new black hole solution obtained for the first time in this work. This solution has a unique feature which may seem strange at first sight. It arises from a coupling of a field of spin 1, with gravitation, which may be the usual Maxwell, or with a contribution of negative energy density, called phantom.

Before beginning the analysis of this new class of phantom black holes, we will present briefly our interest in obtaining and studying such exotic solutions. With the discovery of the acceleration of the universe, various observational programs of study of the evolution of our universe were deployed, including the relationship of the magnitude-versus-redshift type supernovae Ia and the spectrum of anisotropy the cosmic microwave background. These programs promote an accelerated expansion of our universe, which should be dominated by an exotic fluid and should have a negative pressure. Moreover, these observations show that this fluid can be phantom, i.e, with the contribution of negative energy density [7].

As the interest in obtaining these classes has increased, we also found ourselves wanting to analysis of a specific phantom model. We can mention here some recent results in the literature, such as the wormhole solutions [8], the black hole solutions of Einstein-Maxwell-Dilaton theory, [9], the higher-dimensional black holes by Gao and Zhang [10], and the higher-dimensional black branes by Grojean et al [11]. Analyses were also made in algebraic structures of this type of phantom system, as the case of the algebra generated by metrics depending on two temporal coordinates, with $D \geq 5$, which provides phantom fields in $4D$, fulfilled by Hull [12], and Sigma models by Clement et al [13]. Here, we will obtain and study the thermodynamic properties of a solution arising from the coupling of Einstein-Hilbert action with a field of spin 1 and a cosmological constant.

One way to analyse the thermodynamics of black holes that has been widely used is the geometry, for which is defined a thermodynamic phase space, which is metric, and from differential geometry is removed all of the information of the properties of the system. These methods provide us, through the curvature scalar of the space phase, if the system presents thermodynamic interaction, showing were the localisation of the point of the limit of extreme black, and where and when the system passes through a phase transition, and finally, if the system is or no locally or globally stable. These methods have originated in the work of Rao [14], which was subsequently developed by other authors [15]. Other works that have gained attention in the analysis of the thermodynamics of black holes by differential geometry were those of Weinhold [16] and Ruppeiner [17]. We use here so-called Geometrothermodynamics method, however, comparing with the more commonly method used, for verification and interpretation of the obtained results. Geometrothermodynamics appears to be a method which is equivalent or superior in some ways, as for example, due to invariance Legendre transformations and to the consistency of results in the various selected ensembles. Recently, this method has gained some attention by the scientific community [18]. This case, which we will study, is a good test for this method because some difficulties already appeared in the utilization of this method applied to phantom solutions, showing some weakness, or even limitation of the method [19].

This paper is organized as follows. In section 2, we present a summary of the obtaining a new class of solutions studied, arising from the Einstein-(anti)Maxwell theory, and how to obtain their thermodynamic quantities. In section 3, we summarize the method of Geometrothermodynamics. In section 4, we apply the method of Geometrothermodynamics for a new black hole solution and the Reissner-Nordstrom-AdS solution. In section 5, we present our conclusions and perspectives.

2 The field equations and the solutions of phantom Reissner-Nordstrom-(A)dS

The action of this theory is given by

$$S = \int d^4x\sqrt{-g}\left[R + 2\eta F_{\mu\nu}F^{\mu\nu} + 2\Lambda\right] \,, \tag{1}$$

where the first term characterizes of Einstein-Hilbert action, the second is the coupling with the Maxwell field ($\eta = 1$), or a phantom field of spin 1 ($\eta = -1$), and finally the third term is the coupling with the cosmological constant, which can behave as $\Lambda > 0$ (dS) or $\Lambda < 0$ (AdS). The nomenclature known as phantom is due to the fact that the energy density of the field of spin 1 is negative, for $\eta = -1$.

Making the functional variation of the action (1), with respect to the fields A_μ and $g_{\mu\nu}$ (components of the spacetime metric), we get the following field equations

$$\left\{\begin{array}{l} \nabla_\mu F^{\mu\nu} = 0 \,, \\ R_{\mu\nu} = 2\eta\left(\frac{1}{4}g_{\mu\nu}F_{\alpha\beta}F^{\alpha\beta} - F_\mu{}^\alpha F_{\nu\alpha}\right) - \Lambda g_{\mu\nu} \,, \end{array}\right. \tag{2}$$

where we used $R=-4\Lambda$. Considering a static, spherically symmetric spacetime, we can write the line element as

$$ds^2 = e^{2\gamma(r)}dt^2 - e^{2\alpha(r)}dr^2 - e^{2\beta(r)}\left(d\theta^2 + \sin^2\theta d\phi^2\right) \ . \tag{3}$$

The first equation of (2) becomes

$$F^{10} = q\,e^{-(\alpha+\gamma+2\beta)} \ , \ F_{\mu\nu}F^{\mu\nu} = -2q\,e^{-4\beta} \ , \tag{4}$$

where q is a real integration constant. Substituting (4) in the second equation of (2), we get

$$\left\{\begin{array}{lcl} \gamma'' + \gamma'\left(\gamma' - \alpha' + 2\beta'\right) & = & e^{2\alpha}\left(\eta q^2\,e^{-4\beta} - \Lambda\right) \ , \\ \gamma'' + 2\beta'' + \gamma'\left(\gamma' - \alpha'\right) + 2\beta'\left(\beta' - \alpha'\right) & = & e^{2\alpha}\left(\eta q^2\,e^{-4\beta} - \Lambda\right) \ , \\ e^{2(\alpha-\beta)} - \left[\beta'' + \beta'\left(\gamma' - \alpha' + 2\beta'\right)\right] & = & e^{2\alpha}\left(\eta q^2\,e^{-4\beta} + \Lambda\right) \ , \end{array}\right. \tag{5}$$

where $'$ denotes the differentiation with respect to the radial coordinate r. We then choose the quasi-global coordinate system

$$\alpha(r) = -\gamma(r) \ , \ \beta(r) = \ln(r) \ , \tag{6}$$

the second equation of (5) becomes

$$\left(e^{2\gamma}\right)'' + \frac{2}{r}\left(e^{2\gamma}\right)' = 2\left(\eta\frac{q^2}{r^4} - \Lambda\right) \ . \tag{7}$$

Integrating (7), we obtain the following line element and Maxwell field (or phantom) as solutions of the equations of motion of the action (1)

$$\begin{array}{rcl} ds^2 & = & f(r)dt^2 - \left[f(r)\right]^{-1}dr^2 - r^2\left(d\theta^2 + \sin^2\theta d\phi^2\right) \ , \qquad (8) \\ F & = & -\frac{q}{r^2}dr\wedge dt \ , \ f(r) = 1 - \frac{2M}{r} - \frac{\Lambda}{3}r^2 + \eta\frac{q^2}{r^2} \ . \qquad (9) \end{array}$$

Again, the mass M and the constant 1, of $f(r)$, are determined for satisfying the newtonian limit when $\Lambda = 0$, the spacetime being asymptotically that of Minkowski. The solution (8) is that of Reissner-Nordstrom-(A)dS, for $\eta = 1$, and that of anti-Reissner-Nordstrom-(A)dS (phantom), for $\eta = -1$.

The horizons of this solution can be determined by the roots of $f(r)$. We can obtain only two real roots for the equation $f(r) = 0$, which are the external horizon r_+ (events horizon), or internal horizon r_- , where $0 < r_- < r_+$, for $\eta = 1$, and $r_- < 0 < r_+$, for $\eta = -1$. We are interested just to the anti-Reissner-Nordstrom-AdS solution($\Lambda < 0$), and of value of r_+, which can be represented as [35]

$$\begin{array}{rcl} r_+ & = & \frac{1}{2}\left(\sqrt{x} + \sqrt{\frac{6}{\Lambda} - x - \frac{12M}{\Lambda\sqrt{x}}}\right) \ , \qquad (10) \\ x & = & A + B + \frac{2}{\Lambda} \ , \ A = -\sqrt[3]{\frac{2}{y}}\left(\frac{1-4\eta\Lambda q^2}{\Lambda}\right) \ , \ B = -\sqrt[3]{\frac{y}{32}}\left(\frac{3}{\Lambda}\right) \ , \qquad (11) \\ y & = & 2 - 36\Lambda M^2 + 24\eta\Lambda q^2 + \\ & & +\sqrt{(2 - 36\Lambda M^2 + 24\eta\Lambda q^2)^2 - 4(1 - 4\eta\Lambda q^2)^3} \ . \qquad (12) \end{array}$$

The causal structure of the new anti-Reissner-Nordstrom-AdS (anti-RN-AdS) solution, obtained for the first time in this work, is identical to the Schwarzschild solution, but with timelike infinite spacial ($r \to \infty$), as shown in Figure 1. For studying the thermodynamic properties of this solution, it is necessary to express the mass M in terms of the radius of the events horizon r_+ and the charge q as follows. Equating $g_{00} = f(r)$ to zero, we get

$$M = \frac{r_+}{2}\left(1 - \frac{\Lambda}{3} r_+^2 + \eta \frac{q^2}{r_+^2}\right) \; . \tag{13}$$

Now, we are interested to do a geometric analysis, representing the semi-classical effects of the gravitation of the black hole solutions cited previously, i.e, quantizing other fields called matter field, treating classically the gravitational field as the background. Then, we will make the semi-classical thermodynamic of black holes, initiated by Hawking [1] and developed posteriorly by other authors [21].



Figure 1: Carter-Penrose diagram for the new anti-RN-AdS solution.

There exist various forms of obtaining the Hawking temperature, as for example by the Bogoliubov coefficients [22], the metric euclidianization [23], by the energy-momentum tensor [4, 21], by the transmission and reflection coefficients [24, 25], by the analysis of the anomaly term [26] and by the superficial gravity of black hole [27]. Until now, these methods appear to be equivalent [28], then, we propose to use in this work the method of the calculation of the Hawking temperature through the superficial gravity.

The superficial gravity of a black hole is given by the expression [29]:

$$\kappa = \left[\frac{g_{00}'}{2\sqrt{-g_{00}g_{11}}}\right]_{r\,=\,r_+} \; , \tag{14}$$

where r_+ is the radius of the events horizon and the Hawking temperature is related with the superficial gravity by the expression [1, 27]:

$$T = \frac{\kappa}{2\pi} \, . \tag{15}$$

Then, for the case of black hole (8), we found that the superficial gravity (14) is

$$\kappa = \frac{M}{r_+^2} - \frac{\Lambda}{3} r_+ - \eta \frac{q^2}{r_+^2} \, , \tag{16}$$

in which, substituting the mass (13), yields

$$\kappa = \frac{1}{2r_+}\left(1 - \Lambda r_+^2 - \eta\frac{q^2}{r_+^2}\right) . \tag{17}$$

and the Hawking temperature (15), in this case, is :

$$T = \frac{1}{4\pi r_+}\left(1 - \Lambda r_+^2 - \eta\frac{q^2}{r_+^2}\right) . \tag{18}$$

We define the area of the black hole horizon as

$$A = \int_0^{2\pi}\int_0^{\pi} \sqrt{g_{22}g_{33}}\, d\theta d\phi\Big|_{r=r_+} = 4\pi r^2\Big|_{r=r_+} = 4\pi r_+^2 . \tag{19}$$

Then, one can define the entropy of the black hole by [2]

$$S = \frac{1}{4}A = \pi r_+^2 . \tag{20}$$

We also calculate, from (9), the electric potential at the events horizon as

$$A_0 = \int_{+\infty}^{r} F_{10}(r')dr'\Big|_{r=r_+} = \frac{q}{r_+} . \tag{21}$$

Through the equations (13) and (20), we can take the differential of the mass and of the entropy

$$\begin{cases} dM = \frac{1}{2}\left(1 - \Lambda r_+^2 - \eta\frac{q^2}{r_+^2}\right)dr_+ + \eta\frac{q}{r_+}dq , \\ dS = 2\pi r_+ dr_+ , \end{cases} \tag{22}$$

which, with (18) and (21), satisfy the first law of the thermodynamic of black holes [2]

$$dM = TdS + \eta A_0 dq . \tag{23}$$

Note that the first law is generalized for the case anti-RN-AdS, with $\eta = -1$, where the sign of the second term, which is related to the work, changes due to the contribution of negative energy to the system.

We can establish correctly the study of the thermodynamics of the system anti-RN-AdS. From the equation (20) we get $r_+ = \sqrt{S/\pi}$, which inserted in (13) yields

$$M(S, q) = \frac{1}{2}\left(\frac{S}{\pi}\right)^{3/2}\left(\frac{\pi}{S} - \frac{\Lambda}{3} + \eta\pi^2\frac{q^2}{S^2}\right) . \tag{24}$$

From (23), we obtain the following equations of state

$$\left(\frac{\partial M}{\partial S}\right)_q = T , \left(\frac{\partial M}{\partial q}\right)_S = \eta A_0 . \tag{25}$$

By the expression (24), we see that the equations of state (25) are satisfied. Now, we have all the required ingredients for doing an analysis of thermodynamic system of the RN-AdS phantom solution. In next section, we will present a brief summary of Geometrotermodynamic method, which will be applied later.

3 The Geometrothermodynamics

Geometrothermodynamics (GTD) method use the differential geometry tool for representing a physical thermodynamic system. With this, we can construct a mathematical space where it can be defined all the thermodynamic quantities, extensive and intensive, also the postulates and equations of state. Let us consider a $(2n+1)-$dimensional space $\mathbb{T}$, whose the coordinates are thermodynamic potential Φ, extensive variables E^a and intensive variables I^a, where $a = 1, ..., n$. If the space $\mathbb{T}$ possesses a non-degenerated metric $G_{AB}(Z^C)$, where $Z^C = \{\Phi, E^a, I^a\}$, and a Gibbs 1-form $\Theta = d\Phi - \delta_{ab} I^a dE^b$, where δ_{ab} is the Kronecker delta, then if the condition $\Theta \wedge (d\Theta)^n \neq 0$ is satisfied the set $(\mathbb{T}, \Theta, G)$ is called contact riemannian manifold [30]. The space $\mathbb{T}$ is called thermodynamic phase space.

One can also define a n-dimensional subspace $\mathbb{E} \subset \mathbb{T}$ with extensive coordinates E^a and a map $\varphi : \mathbb{E} \to \mathbb{T}$, with $\Phi \equiv \Phi(E^a)$, such that

$$\varphi^*(\Theta) \equiv 0 \Rightarrow \left\{ \begin{array}{l} d\Phi = \delta_{ab} I^a dE^b \ , \\ \frac{\partial \Phi}{\partial E^a} = \delta_{ab} I^b \ . \end{array} \right. \tag{26}$$

Then, we call the space $\mathbb{E}$ by the space of equilibrium thermodynamic states, and first equation of (26) by "first law of thermodynamics", and the second equation by the condition for thermodynamic equilibrium. We impose "the second law of thermodynamics" as a required condition:

$$\pm \frac{\partial^2 \Phi}{\partial E^a \partial E^b} \geq 0 \ , \tag{27}$$

where the sign $(\pm)$ depends on the chosen thermodynamic potential. In the mass case we get the sign $(+)$ and for the entropy case we get $(-)$. This condition is known as convexity of the thermodynamic potential. The thermodynamic potential is defined in such a way to satisfy the condition of homogeneity $\Phi(\alpha E^a) = \alpha^\beta \Phi(E^a)$. Differentiating with respect to α, using the second equation in (26) and taking $\alpha = 1$, we obtain

$$\beta \Phi(E^a) = \delta_{ab} I^a E^b \ . \tag{28}$$

Differentiating (28) and using the first equation of (26), we get

$$(1-\beta)\delta_{ab} I^a dE^b + \delta_{ab} E^a dI^b = 0 \ . \tag{29}$$

For $\beta = 1$, we re-obtain the Euler identity in (28), and the relation of Gibbs-Duhem in (42). The pullback $\varphi^* : T^*(\mathbb{T}) \otimes T^*(\mathbb{T}) \to T^*(\mathbb{E}) \otimes T^*(\mathbb{E})$[4], induces a metric in $\mathbb{E}$, such that $\varphi^*(G) = g$.

Hernando Quevedo has improved probable metric G for the GTD, such that for the case of black holes, it can be written as follows [5]

$$dL^2 = G_{AB} dZ^A dZ^B = \Theta^2 + \left(\delta_{ab} E^a I^b\right) \left(\eta_{ab} dE^a dI^b\right) \ , \tag{30}$$

---

[4] Where $T^*(\mathbb{T})$ and $T^*(\mathbb{E})$ represent the tangent space to $\mathbb{T}$ and $\mathbb{E}$ respectively.

where $\eta_{ab} = \{\pm 1, 1, ..., 1\}$, which the second order phase transition leads to $\eta_{ab} = \{-1, 1, ..., 1\}$. Pullback φ^* induces a metric in $\mathbb{E}$:

$$\begin{aligned} dl^2 = g_{ab} dE^a dE^b &= \frac{\partial Z^A}{\partial E^a} \frac{\partial Z^B}{\partial E^b} G_{AB} dE^a dE^b , \\ &= \left(E^c \frac{\partial \Phi}{\partial E^c} \right) \left(\eta_{ad} \delta^{di} \frac{\partial^2 \Phi}{\partial E^i E^b} \right) dE^a dE^b \ . \end{aligned} \tag{31}$$

Since the thermodynamic system does not depend on the choice of the thermodynamic potential and it is invariant by the Legendre transformations, we found that the metrics (30) and (31) are defined in such a way to invariants by the Legendre transformations as follows:

$$\Phi = \widehat{\Phi} - \delta_{ab} \widehat{E}^a \widehat{I}^b \ , \ E^a = -\widehat{I}^a \ , \ I^b = \widehat{E}^b \ . \tag{32}$$

The n-dimensional space $\mathbb{E}$, whose the metric is g_{ab}, is used for interpreting the thermodynamic interactions, phase transitions and fluctuations or stability of the thermodynamic system. Through the metric (31), we can calculate the scalar of curvature R, which provides two interpretations: if it exists thermodynamic interaction and phase transition, and at which point of the space of thermodynamic equilibrium states they occur.

We have seen that a metric of the space of thermodynamic equilibrium states $\mathbb{E}$ can be obtained by the pullback of the metric of the contact riemannian space. By the definition of the line element (31), of the space $\mathbb{E}$, one can define the distribution of probability in the interval $E^a + dE^a$ for finding the physical state with the extensive variable E^a, by

$$P(E^a) = \frac{\sqrt{\det [g_{ab}]}}{(2\pi)^{\frac{n}{2}}} \exp \left[\frac{1}{2} g_{ab} dE^a dE^b \right] \ , \tag{33}$$

where $P(E^a)$ satisfies

$$\int \prod_{a=1}^{n} dE^a P(E^a) = 1 \ . \tag{34}$$

Deriving (33) with respect to V^{-1}, where V is the volume of the system, the expression for the second fluctuations can be obtained (in the thermodynamic limit $V \to \infty$) [31]

$$\langle \Delta E^a \Delta E^b \rangle = -g^{ab} \ , \tag{35}$$

where $\Delta E^a = E^a - E^a_{(0)}$ and g^{ab} is the inverse of g_{ab}. The unities must be adjusted with a realistic analysis. In the case where the fluctuations are real and small the system is stable.

Another criterion used for determining the stability is through the following objects

$$p_1^{(1)} \ = \ g_{11} > 0 \ , \ p_1^{(2)} = g_{22} > 0 \ , \ ... \ , \ p_1^{(n)} = g_{nn} > 0 \ , \tag{36}$$

$$p_2^{(1)} = \begin{vmatrix} g_{11} & g_{12} \\ g_{12} & g_{22} \end{vmatrix} > 0 \ , \ p_2^{(2)} = \begin{vmatrix} g_{22} & g_{23} \\ g_{23} & g_{33} \end{vmatrix} > 0 \ , \tag{37}$$

$$p_3^{(1)} = \begin{vmatrix} g_{11} & g_{12} & g_{13} \\ g_{12} & g_{22} & g_{23} \\ g_{13} & g_{23} & g_{33} \end{vmatrix} > 0 \ , \ p_n = \det\left[g_{ab}\right] > 0 \ . \tag{38}$$

The positive (negative) sign of p_n, which depends on the choice of the thermodynamic potential, gives us the notion of the local stability (instability) of the thermodynamic system and the fulfilment of [32]

$$p_i > 0 \ , \ i = 1, \ldots, n \ , \tag{39}$$

informs us that the system is globally stable.

Since we are not studying rotating black holes, another mechanism for determining the global stability (instability) is the Helmholtz free energy. In thermodynamic variables of black holes, the Helmholtz free energy is a Legendre transformation of the mass (energy) $M(S, q)$:

$$F(T, q) = M(S, q) - TS \ . \tag{40}$$

When

$$F(E^a) < 0 \ , \ \forall E^a \ (E^a \in I(E^a)) \ , \tag{41}$$

where $I(E^a)$ is an interval and E^a are the extensive variables, the thermodynamic system is globally stable. In the usual case of the thermodynamics the Helmholtz free energy is given by $F(T, V) = U - TS$.

We also can define the Gibbs potential as

$$G(T, A_0) = M(S, q) - TS - \eta A_0 q \ , \tag{42}$$

and the global stability is given by

$$G(E^a) < 0 \ , \ \forall E^a \ (E^a \in I(E^a)) \ . \tag{43}$$

We introduced the sign of η in (42) for compensating the contribution of the thermodynamic work. Then, we will use the Gibbs potential for determining the global stability of the thermodynamic system. In next section, we will explain these methods of study of the thermodynamic properties of the solutions RN-Ads and anti-RN-Ads.

4 Termodynamics of phantom Reissner-Nordstrom-AdS black hole

4.1 Geometrothemodynamics Application

We start defining the thermodynamic variables of the system. For the cases of black holes coming from the Einstein-Maxwell theory, we always will get a solution with two physical parameters, the mass M and the charge q. Other elements which could be defined from these two parameters are the entropy S, the temperature T and the electric potential scalar A_0. The contact riemannian manifold $\mathbb{T}$ is, in this case, a 5-dimensional space and the space of thermodynamic equilibrium states $\mathbb{E}$ is a 2-dimensional submanifold.

The description of the thermodynamics is that of the mass representation $M(S,q)$ in (24), as thermodynamic potential Φ, defined in the previous section. The extensive variables are the entropy S and the charge q, which was represented by the coordinates E^a. The intensives variables are the temperature T and the electric potential A_0, which was represented by the coordinates I^a.

Then, we have the coordinates of the thermodynamic phase space $\mathbb{T}$ as $Z^A = \{M(S,q), S, q, T, A_0\}$, and the Gibbs 1-form is given by [5]

$$\Theta_M = dM - TdS - \eta A_0 dq \; , \tag{44}$$

such that we have $\varphi^*(\Theta_S) = 0$, resulting first law of the thermodynamics of black holes $dM = TdS + \eta A_0 dq$ ($\eta = \pm 1$).

The line element (30), of the space $\mathbb{T}$, for a second order phase transition is given by

$$\begin{aligned} dL^2 &= (dM - TdS - \eta A_0 dq)^2 + (TS + \eta A_0 q) \times \\ &\quad \times \left[-dSdT + d\left(\eta A_0\right) dq\right] \; . \end{aligned} \tag{45}$$

The first law, second law and the equations of state at the equilibrium are given by

$$d\Phi = \delta_{ab} I^a dE^b \rightarrow dM = TdS + \eta A_0 dq \; , \tag{46}$$

$$\frac{\partial^2 M}{\partial S^2} , \frac{\partial^2 M}{\partial S \partial q} , \frac{\partial^2 M}{\partial q^2} \leqslant 0 \; , \tag{47}$$

$$\frac{\partial \Phi}{\partial E^a} = \delta_{ab} I^b \rightarrow \frac{\partial M}{\partial S} = T \, , \; \frac{\partial M}{\partial q} = \eta A_0 \; . \tag{48}$$

Now, we must specify the solution anti-RN-AdS (8). The line element (31) of the space of thermodynamic equilibrium states, taking into account (24), is given by

$$dl^2 = \left(S \frac{\partial M}{\partial S} + q \frac{\partial M}{\partial q} \right) \left(- \frac{\partial^2 M}{\partial S^2} dS^2 + \frac{\partial^2 M}{\partial q^2} dq^2 \right) \; , \tag{49}$$

$$= g_{SS} dS^2 + g_{qq} dq^2 , \tag{50}$$

$$g_{SS} = \left(\frac{\pi S - \Lambda S^2 + 3\eta \pi^2 q^2}{4\pi S} \right) \left(\frac{\Lambda S^2 + \pi S - 3\eta \pi^2 q^2}{8\pi^2 S^2} \right) \; , \tag{51}$$

$$g_{qq} = \eta \left(\frac{\pi S - \Lambda S^2 + 3\eta \pi^2 q^2}{4\pi S} \right) \; . \tag{52}$$

---

[5] This expression comes from the first law of thermodynamics (23).

We note here that the phantom contribution may change the signature of the space metric $\mathbb{E}$. For calculating the scalar of curvature associated to the metric (50), we can use the formula of a scalar of a two dimensional space:

$$\begin{aligned} R(M,q) &= -\frac{1}{\sqrt{|det[g]|}}\left[\partial_q\left(\frac{\partial_q g_{MM}-\partial_M g_{Mq}}{\sqrt{|det[g]|}}\right)+\partial_M\left(\frac{\partial_M g_{qq}-\partial_q g_{Mq}}{\sqrt{|det[g]|}}\right)\right] \\ &\quad -\frac{det[H_S]}{2\left(det[g]\right)^2}\,, \qquad (53) \\ H_S &= \begin{pmatrix} g_{MM} & g_{Mq} & g_{qq} \\ \partial_M g_{MM} & \partial_M g_{Mq} & \partial_M g_{qq} \\ \partial_q g_{MM} & \partial_q g_{Mq} & \partial_q g_{qq} \end{pmatrix}\,. \qquad (54) \end{aligned}$$

The scalar (53) is given by

$$\begin{aligned} R(S,q) &= -16\pi^3 S^2\Big(5\Lambda^3 S^5+\pi\Lambda^2 S^4-2\pi^2\Lambda S^3+18\eta\pi^4 S q^2+15\eta\Lambda\pi^3 S^2 q^2+ \\ &+ 6\eta\Lambda^2\pi^2 S^3 q^2-18\pi^5 q^4-63\Lambda\pi^4 S\Big)\Big/\left(\Lambda S^2+\pi S-3\eta\pi^2 q^2\right)^2\times \\ &\times \left(\Lambda S^2-\pi S-3\eta\pi^2 q^2\right)^3\,. \qquad (55) \end{aligned}$$

By the use of a mathematical software, we found the roots of the scalar of curvature $R(M,q)$ in (55) for the values of the entropy $S_1 = 0.0305608$ and $S_2 = 2.441077$ when $\eta = -\Lambda = 1$ and $q = 0.1$; and $S_3 = 0.864182$ and $S_4 = 2.22539$ when $\eta = \Lambda = -1$ and $q = 0.1$. The points where the scalar $R(M,q)$ diverges are given by the values of the entropy $S_5 = -(\pi/2\Lambda)(1+\sqrt{1+12\eta\Lambda q^2})$, $S_6 = -(\pi/2\Lambda)(-1+\sqrt{1+12\eta\Lambda q^2})$ and $S_7 = -(\pi/2\Lambda)(1-\sqrt{1+12\eta\Lambda q^2})$. All the other points are negative or complex values for the entropy and have been rejected.

For confirming the consistency of our analysis, we can calculate in the usual thermodynamic way the specific heat by the expression

$$C_q = \left(\frac{\partial M}{\partial T}\right)_q = \left(\frac{\partial M}{\partial S}\right)_q \Big/ \left(\frac{\partial^2 M}{\partial S^2}\right)_q\,, \qquad (56)$$

which leads to

$$C_q = 2S\frac{\left(-\pi S+\Lambda S^2+\eta\pi^2 q^2\right)}{\left(\pi S+\Lambda S^2-3\eta\pi^2 q^2\right)}\,. \qquad (57)$$

The roots of (57) give the values where the black hole is extreme, which are $S_8 = -(\pi/2\Lambda)(-1+\sqrt{1-4\eta\Lambda q^2})$ and $S_9 = (\pi/2\Lambda)(1+\sqrt{1-4\eta\Lambda q^2}) < 0$, where the last value has been rejected since it is negative ($\Lambda < 0$). The values S_8 and S_9 can be obtained directly setting the temperature (18) equal to zero and substituting $r_+ = \sqrt{S/\pi}$. The points where the specific heat diverges, i.e, the points of phase transitions, are S_5 and S_7 given previously. More particularly here, where only the value of S_5 is positive for the case anti-RN-AdS, i.e, this case possesses just one point of phase transition. This result is in agreement with that known in the literature, which commonly is demonstrated for the specific heat in terms of the events horizon [33], which in our case is a direct substitution of (20).

We present the graphics of the scalar of curvature (55) and the specific heat (57) in function of the entropy, for the fix value of the electric charge $q = 0.25$, in the Figures 2 and 3. For the case RN-AdS, the unique root for the specific heat is $S_8(q = 0.1) = 0.0311079$, which differs from the two values which make the scalar of curvature vanishing, $S_1(q = 0.1) = 0.0305608$ and $S_2(q = 0.1) = 2.441077$. In the case anti-RN-AdS, there is any root for which the specific heat vanishes, but there are two roots which the scalar of curvature vanishes, $S_3(q = 0.1) = 0.864182$ and $S_4(q = 0.1) = 2.22539$. Thus, Geometrothermodynamics method does not provide the same result for the analysis done by the specific heat of the black hole, both for the RN-AdS and for the anti-RN-AdS cases. However, again, the analysis performed by the Geometrothermodynamics method appears to be inefficient, as it has been shown in [19]. The graphics of the scalar of curvature and the specific heat, for this case, are presented in Figures 4 and 5.



Figure 2: Graph of the scalar of curvature in function of the entropy S, of the RN-AdS case, for the electric charge $q = 0.25$ and the cosmological constant $\Lambda = -1$.



Figure 3: Graph of the specific heat in function of the entropy S, of the RN-AdS case, for the electric charge $q = 0.25$ and the cosmological constant $\Lambda = -1$.

4.2 Local and global stability

The local stability of a thermodynamic system can be studied commonly by the specific heat. But we can also study the components of the metric of the space



Figure 4: Graph of the scalar of curvature in function of the entropy S, of the anti-RN-AdS case, for the electric charge $q = 0.25$ and the cosmological constant $\Lambda = -1$.



Figure 5: Graph of the specific heat in function of the entropy S, of the anti-RN-AdS case, for the electric charge $q = 0.25$ and the cosmological constant $\Lambda = -1$.

of thermodynamic equilibrium states $\mathbb{E}$ or the Hessianmass. About the global instability of the system, it can be determined by the analysis of components of the metric $\mathbb{E}$ and all possible determinants of them, by the Helmholtz free energy or by the Gibbs potential. We will study here the local and global stability of the class of the black holes solutions of the type RN-AdS and anti-RN-AdS.

Let us start calculating the Hessian of the mass (24). The Hessianmatrix of the mass is defined by

$$H_S = \begin{pmatrix} \frac{\partial^2 M}{\partial S^2} & \frac{\partial^2 M}{\partial S \partial q} \\ \frac{\partial^2 M}{\partial S \partial q} & \frac{\partial^2 M}{\partial q^2} \end{pmatrix} , \tag{58}$$

which for (24) yields

$$H_S = \begin{pmatrix} \frac{3\eta\pi^2 q^2 - \pi S - \Lambda S^2}{8\pi^{3/2} S^{5/2}} & -\eta\frac{q\sqrt{\pi}}{2S^{3/2}} \\ -\eta\frac{q\sqrt{\pi}}{2S^{3/2}} & \eta\sqrt{\frac{\pi}{S}} \end{pmatrix} . \tag{59}$$

The components of the Hessianmatrix assume clearly the positive and negative values, which implies the local instability for the two solutions, RN-AdS and anti-RN-AdS.

We can also analyse the specific heat (57), but to do this, let us first study the temperature of the black hole (18). The temperature can be written in terms of the entropy as follows

$$T(S, q) = \frac{\pi S - \Lambda S^2 - \eta\pi^2 q^2}{4(\pi S)^{3/2}} . \tag{60}$$

In the RN-AdS case the temperature is positive for $S > S_e$, where the value for to extreme case is given by $S_e = (-\pi/2\Lambda)(-1 + \sqrt{1 - 4\Lambda q^2})$, zero for $S = S_e$ and negative for $S < S_e$ (unstable), which is in accordance with [34, 35]. For the anti-RN-AdS case, the temperature is always positive for $S > 0$, then there isn't the extreme case, as we previously mentioned. The graphics of the Figures 6 and 7 are of temperature in function of the entropy for the two cases. In the RN-AdS case, it is clear that $T = 0$ for S_e, passing later by the maximum and decreasing slowly. For the anti-RN-AdS case, the temperature starts very great, decreasing until a minimum and later increases more slowly.



Figure 6: Graph of the temperature in function of the entropy S, of the RN-AdS case, for the electric charge $q = 0.1$ and the cosmological constant $\Lambda = -1$.



Figure 7: Graph of the temperature in function of the entropy S, of the anti-RN-AdS case, for the electric charge $q = 0.1$ and the cosmological constant $\Lambda = -1$.

Now, using (60) we can write the specific heat (57) as follows

$$C_q = \frac{8T(S,q)\pi^{3/2}S^{5/2}}{3\eta\pi^2 q^2 - \pi S - \Lambda S^2} \,. \tag{61}$$

Analysing the sign of the specific heat for the RN-AdS case, we have a locally unstable phase where $C_q < 0$, for $S \in (S_7, S_5)$, where $S_{5,7} = -(\pi/2\Lambda)(1 \pm \sqrt{1 + 12\eta\Lambda q^2})$ and $S_e < S_7$. The black hole is extreme in $S = S_e$ and one has a local stable phase where $C_q > 0$, for $S \in (S_e, S_7)$ or $S > S_5$, with $S_e < S_7$, according to the graphic of the Figure 3. This also provides a value of the extreme charge, to the RN-AdS case, which is given by $q_e = 1/2\sqrt{3} \approx 0.28867513459$ [34]. The anti-RN-AdS case, the graphic of the Figure 5 shows that here is not extreme case and the system is locally stable for the condition $S > S_5$, since $Cq < 0$ for $0 < S < S_5$.

We can also study the local and global stability by the metric components of the space of thermodynamic equilibrium states $\mathbb{E}$ in (49), as shown in the section 3. The sign of the metric components (49) for the RN-AdS case are given by $g_{SS} > 0$ for $S \in (S_7, S_5)$, $g_{qq} > 0$ for $S > 0$ and the determinant of the metric $g > 0$ for $S \in (S_7, S_5)$. Collecting the results, we observe that the system is local and globally stable if $S \in (S_7, S_5)$. This is not in agreement with the analysis made by the specific heat. But for the anti-RN-AdS case, the components g_{SS}, g_{qq} and the determinant g are always positive in disjunct interval. Then there does not exist interval for which the system can be locally or globally stable, contradicting again the result by the specific heat.

Now, we will study the global stability of these solutions. We can rewrite the mass and the electric potential in terms of the entropy and of the electric charge as follows

$$M = \frac{-\Lambda S^2 + 3\pi S + 3\eta\pi^2 q^2}{6\pi^{3/2}\sqrt{S}} \ , \ A_0 = q\sqrt{\frac{\pi}{S}} \ . \tag{62}$$

Our analysis starts with the grand-canonical ensemble. Using (60) and (62), one can calculate the Gibbs potential (42)

$$G = \frac{\Lambda S^2 + 3\pi S - 3\eta\pi^2 q^2}{12\pi^{3/2}\sqrt{S}} \ . \tag{63}$$

Equating (63) to zero, one has

$$S_{G1,G2} = \left(-\frac{\pi}{2\Lambda}\right)\left(3 \mp \sqrt{9 + 12\eta\Lambda q^2}\right) \ . \tag{64}$$

In the RN-AdS case, by the Gibbs potential, we get $G < 0$, which is a stable system for $S \in (0, S_{G1}) \cup (S_{G2}, +\infty)$, which is in accordance with [35]. In the anti-RN-AdS case, one gets $G < 0$, which is a stable system only when $S > S_{G2}$. The graphs of the Gibbs potential for the two cases are presented in the Figures 8 and 9.



Figure 8: Graph of the Gibbs potential, to the RN-AdS case, in function of the entropy S for the electric charge $q = 0.1$ and the cosmological constant $\Lambda = -1$.

We can now analyse the canonical ensemble. The Helmholtz free energy (40), from (60) and (62), is given by

$$F = \frac{\Lambda S^2 + 3\pi S + 9\eta\pi^2 q^2}{12\pi^{3/2}\sqrt{S}} \ , \tag{65}$$



Figure 9: Graph of the Gibbs potential, to the anti-RN-AdS case, in function of the entropy S for the electric charge $q = 0.1$ and the cosmological constant $\Lambda = -1$.

which equating to zero yields

$$S_{F1,F2} = \left(-\frac{3\pi}{2\Lambda}\right)\left(1 \pm \sqrt{1 - 4\eta\Lambda q^2}\right) \ . \tag{66}$$

Through the canonical ensemble it seems that we get the inversion of the cases. In the RN-AdS case the system is stable, i.e, $F < 0$ for $S > S_{F1}$. In the anti-RN-AdS case the system is stable for $S \in (0, S_{F2}) \cup (S_{F1}, +\infty)$. The graph of the Helmholtz free energy, for the two cases, are represented in the Figures 10 and 11.



Figure 10: Graph of the Helmholtz free energy, of the RN-AdS case, in function of the entropy S for the value of the electric charge $q = 0.1$ and the cosmological constant $\Lambda = -1$.



Figure 11: Graph of the Helmholtz free energy, of the anti-RN-AdS case, in function of the entropy S for the values of the electric charge $q = 0.1$ and the cosmological constant $\Lambda = -1$.

5 Conclusions

We got a new solution of Einstein-anti-Maxwell theory with cosmological constant, which is called anti-RN-(A)dS solution. We analysed the basic properties of this solution, such as the horizons. Later, we focused our attention to the thermodynamics of the anti-RN-AdS solution.

We made a summary of the Geometrothermodynamics method. We applied the method to the solutions of types RN-AdS and anti-RN-AdS, as well as the more usual method of black holes thermodynamics. Considering the mass like the thermodynamic potential, we obtained the scalar curvature (55) of the space $\mathbb{E}$. This scalar possesses two roots and two points of phase transition in each case. But these roots are not obtained by the specific heat method. For the RN-AdS case, the unique root for the specific heat is $S_8(q = 0.1) = 0.0311079$, which differs from the two values which cancel the scalar curvature $S_1(q = 0.1) = 0.0305608$ and $S_2(q = 0.1) = 2.441077$. In anti-RN-AdS case, there is any root for the specific heat, but there is two roots of the curvature, $S_3(q = 0.1) = 0.864182$ and $S_4(q = 0.1) = 2.22539$. So, the Geometrothermodynamics method does not give us the same result of the analysis made by the specific heat of the black hole, in both cases, i.e, RN-AdS and anti-RN-AdS. Again, the analysis made by the Geometrothermodynamics method has been ineffective in some way, as in [19].

Then we analyse the local and global stability of these solutions, through the Hessian of the mass, of the components and the determinant of the metric of the space of equilibrium thermodynamic states $\mathbb{E}$, the specific heat, the Gibbs potential and Helmholtz free energy. The local stability can not be established by the Hessian of the mass. For a given interval of real values of the entropy, the local stability can be established for the two solutions, using the specific heat. Using the components and determinant of the metric in $\mathbb{E}$ we do not obtain a result consistent with to calculus by the specific heat. Finally, through the grand-canonical and canonical ensembles, we establish the global stability of the thermodynamic system of the two solutions, RN-AdS and anti-RN-AdS. These ensembles seem to produce results symmetric with each other.

Then we showed the consistent construction of the thermodynamics of the anti-RN-AdS solution and its global stability. This shows clearly that this solution is physically stable. We hope that this analysis of the thermodynamics system made in this paper leads to a constraint to the spacetime geometry stability of this solution.

About the Geometrothermodynamics method, we hope that establishing a new metric of the thermodynamic space $\mathbb{T}$, we will can interpret the results more consistently. We believe that the introduction *ad hoc* of the Lorentzian signature (30), it must have a more physical justification. In fact, when we introduce this metric we are to impose the result that we expect for to thermodynamic system. For example, when we introduced the Lorentzian metric, we imposed to the thermodynamic system a possible second order phase transition. It will be necessary a greater understanding of the geometry of contact riemannian manifold, so that we can introduce a metric arising from previously established physical requirements, but which may offer posteriorly no known results.

References

[1] S. Hawking, Commun. Math. Phys. **43**, 199 (1975).

[2] J. M. Bardeen, B. Carter and S. W. Hawking, Commun. Math. Phys., 31, 161 (1973).

[3] C. O. Lousto, Nucl.Phys. B **410**: 155-172 (1993); Erratum-ibid. **449**: 433 (1995).

[4] Proc.Roy.Soc.Lond. A **353**: 499-521 (1977).

[5] H. Quevedo, A. Sanchez, S. Taj and A. Vazquez, Gen. Rel. Grav. **43**:1153-1165 (2011).

[6] Jorma Louko and Stephen N. Winters-Hilt, Phys. Rev. D **54**:2647-2663 (1996).

[7] S. Hannestad, Int. J. Mod. Phys. A **21**, 1938 (2006); J. Dunkley et al., Astrophys. J. Suppl. Ser. **180**, 306 (2009).

[8] K.A. Bronnikov, M.S. Chernakova, J.C. Fabris, N. Pinto-Neto and M.E. Rodrigues, Int.J.Mod.Phys.D **17**:25-42 (2008).

[9] G.W. Gibbons and D. A. Rasheed, Nucl. Phys. B **476**, 515 (1996).

[10] C. J. Gao and S. N. Zhang, arXiv:hep-th/0604114.

[11] C. Grojean, F. Quevedo, G. Tasinato, and I. Zavala, J. High Energy Phys. **08** (2001) 005.

[12] C. M. Hull, JHEP **9807** (1998) 021.

[13] Mustapha Azreg-Anou, Grard Clment, Jlio C. Fabris, Manuel E. Rodrigues, Phys.Rev.D **83**:124001 (2011).

[14] C. R. Rao, Bull. Calcutta Math. Soc. **37**, 81 (1945).

[15] S. Amari, Differential-Geometrical methods in Statistics (Springer-Verlag, Berlin, 1985).

[16] F. Weinhold, J. Chem. Phys. **63**, 2479, 2484, 2488, 2496 (1975); **65**, 558 (1976).

[17] G. Ruppeiner, Phys. Rev. A **20**, 1608 (1979); G. Ruppeiner, Rev. Mod. Phys. **67**, 605 (1995); **68**, 313 (1996).

[18] Danny Birmingham, Susan Mokhtari, Phys.Lett.B **697**:80-84 (2011), arXiv:1011.6654v1 [hep-th]; Hernando Quevedo, Maria N. Quevedo, arXiv:1111.5056v1 [math-ph]; Hernando Quevedo, Alberto Sanchez, Phys.Rev.D **79**:024012 (2009), arXiv:0811.2524v1 [gr-qc]; Alexis Larranaga, Alejandro Cardenas, arXiv:1108.2205v1 [gr-qc]; Hernando Quevedo, Alberto Sanchez, Safia Taj, Alejandro Vazquez, arXiv:1101.4494v1 [hep-th]; Alexis Larranaga, Sindy Mojica, Brazilian Journal of Physics **41**: 154-158 (2011), arXiv:1012.2070v1 [gr-qc]; Hernando Quevedo, Alberto Sanchez, JHEP **0809**:034 (2008), arXiv:0805.3003v2 [hep-th]; Hernando Quevedo, Alberto Sanchez, Phys. Rev. D **79**, 087504 (2009), arXiv:0902.4488v2 [gr-qc]; Safia Taj, Hernando Quevedo, arXiv:1104.3195v1 [math-ph]; W. Janke, D. A. Johnston, R. Kenna, J. Phys. A **43**:425206, 2010, arXiv:1005.3392v2 [hep-th]; Lszl rpd Gergely, Narit Pidokrajt, Sergei Winitzki, Eur.Phys.J.C **71**:1569 (2011), arXiv:0811.1548v3 [gr-qc]; Hernando Quevedo, J.Math.Phys. **48** (2007) 013506, arXiv:physics/0604164v2 [physics.chem-ph]; Hernando Quevedo, Alberto Sanchez, Alejandro Vazquez, arXiv:0811.0222v1 [math-ph]; H. Quevedo, A. Sanchez, S. Taj, A. Vazquez, Gen.Rel.Grav. **43**:1153-1165,2011, arXiv:1010.5599v1 [gr-qc]; Hernando Quevedo, Alejandro Vazquez, AIP Conf.Proc. **977**:165-172 (2008), arXiv:0712.0868v1 [math-ph]; M. Akbar, H. Quevedo, K. Saifullah, A. Sanchez, S. Taj, Phys.Rev.D **83**: 084031 (2011), arXiv:1101.2722v1 [gr-qc]; H. Quevedo, A. Sanchez, A. Vazquez, arXiv:0805.4819v5 [hep-th]; Alejandro Vazquez, Hernando Quevedo, Alberto Sanchez, J.Geom.Phys. **60**: 1942-1949 (2010), arXiv:1101.3359v1 [math-ph]; J.L. Alvarez, H. Quevedo, A. Sanchez, Phys.Rev.D **77**: 084004 (2008),arXiv:0801.2279v1 [gr-qc].

[19] Manuel E. Rodrigues and Zui A. A. Oporto, Thermodynamics of phantom black holes in Einstein-Maxwell-Dilaton theory, arXiv:1201.5337 [gr-qc].

[20] Ratna Koley, Joydip Mitra, Supratik Pal and Soumitra SenGupta, A critical analysis of thermodynamic properties of braneworld black holes in anti-de Sitter spacetime, arXiv:0910.5096v1 [hep-th].

[21] N. D. Birrell and P. C. W. Davies, Quantum fields in curved space, Cambridge University Press, 1982.

[22] L. H. Ford. arXiv: gr-qc/9707062.

[23] G. W. Gibbons and S. Hawking, Phys. Rev. D **15**: 2752-2756 (1977).

[24] G. Clement, J. C. Fabris and G. T. Marques, Phys. Lett. B **651**: 54-57 (2007).

[25] Panagiota Kanti and John March-Russell, Phys.Rev.D **66**: 024023 (2002); Wontae Kim and John J. Oh, J.Korean Phys.Soc. **52**: 986 (2008); Kazuo Ghoroku, Arne L. Larsen, Phys.Lett. B **328**: 28-35 (1994).

[26] S.P. Robinson and F. Wilczek, Phys. Rev. Lett. **95**: 011303 (2005).

[27] T. Jacobson and G. Kang, Class.Quant.Grav. **10**:L201-L206 (1993); arXiv: gr-qc/9307002.

[28] Glauber Tadaiesky Marques and Manuel E. Rodrigues, Eur. Phys. J. C 72: 1891 (2012), arXiv:1110.0079v2 [gr-qc].

[29] R. M. Wald, General Relativity, Univ. of Chicago Press, 1984, Chicago.

[30] R. Hermann, Geometry, physics and systems (Marcel Dekker, New York, 1973); G. Hernandez and E. A. Lacomba, Contact Riemannian geometry and thermodynamics, Diff. Geom. and Appl. 8, 205 (1998).

[31] George Ruppeiner, Phys. Rev. D **75**, 024037 (2007); Rev. Mod. Phys. **67**: 605-659 (1995), Erratum-ibid. **68**: 313-313 (1996).

[32] Stefano Bellucci and Bhupendra Nath Tiwari, arXiv:1103.2064v1 [hep-th].

[33] Chao Niu, Yu Tian and Xiaoning Wu, Phys.Rev. D **85**: 024017 (2012).

[34] Rabin Banerjee, Sumit Ghosh and Dibakar Roychowdhury, Phys.Lett. B **696**: 156-162 (2011).

[35] Anurag Sahay, Tapobrata Sarkar and Gautam Sengupta, JHEP **1007**: 082 (2010).